

Chapter 28: Delta Functions

28.1 a. $|I| = |m[v_{\text{after}} - v_{\text{before}}]| = |0.145[45 - 0]| = 6.525 \text{ (kg}\cdot\text{meter/sec)}$.

28.2 a. Solving the equation $I = m[v_{\text{after}} - v_{\text{before}}]$ for v_{after} we get

$$v_{\text{after}} = \frac{I}{m} + v_{\text{before}} \text{ .}$$

In particular, assuming $m = 2$ and $v_{\text{before}} = -10$:

If $I = 60$, then $v_{\text{after}} = \frac{60}{2} - 10 = 20 \text{ (meter/sec)}$.

If $I = 100$, then $v_{\text{after}} = \frac{100}{2} - 10 = 40 \text{ (meter/sec)}$.

If $I = 20$, then $v_{\text{after}} = \frac{20}{2} - 10 = 0 \text{ (meter/sec)}$.

28.2 c. Solving the equation $I = m[v_{\text{after}} - v_{\text{before}}]$ for m we get

$$m = \frac{I}{v_{\text{after}} - v_{\text{before}}} \text{ .}$$

In particular, assuming $I = 30$ and $v_{\text{before}} = -10$:

If $v_{\text{before}} = -10$ and $v_{\text{after}} = 50$, then $m = \frac{30}{50 - (-10)} = \frac{1}{2} \text{ (kg)}$.

If $v_{\text{before}} = 0$ and $v_{\text{after}} = 15$, then $m = \frac{30}{15 - 0} = 2 \text{ (kg)}$.

28.3 a. $\int_0^{\infty} t^2 \delta_4(t) dt = 4^2 = 16$.

28.3 c. $\int_0^{\infty} \cos(t) \delta(t) dt = \int_0^{\infty} \cos(t) \delta_0(t) dt = \cos(0) = 1$.

28.3 e. $\int_0^{\infty} t^2 \text{rect}_{(1,4)}(t) \delta_3(t) dt = 3^2 \text{rect}_{(1,4)}(3) = 9 \cdot 1 = 9$.

28.5. Keeping in mind theorem 28.1, and the fact that $\alpha \geq 0$ and $t > 0$, we have

$$\begin{aligned} g * \delta_{\alpha}(t) &= \delta_{\alpha} * g(t) = \int_0^t \delta_{\alpha}(x) g(t-x) dx \\ &= \int_0^{\infty} \delta_{\alpha}(x) g(t-x) \text{rect}_{(0,t)}(x) dx \\ &= g(t-\alpha) \text{rect}_{(0,t)}(\alpha) \\ &= g(t-\alpha) \cdot \begin{cases} 1 & \text{if } 0 < \alpha < t \\ 0 & \text{if } t < \alpha \end{cases} \\ &= g(t-\alpha) \cdot \begin{cases} 0 & \text{if } t < \alpha \\ 1 & \text{if } \alpha < t \end{cases} = g(t-\alpha) \text{step}_{\alpha}(t) \text{ .} \end{aligned}$$

28.6 a. $\mathcal{L}[y']|_s = \mathcal{L}[3\delta_2(t)]|_s$

$\hookrightarrow sY(s) - 0 = 3\mathcal{L}[\delta_2(t)]|_s = 3e^{-2t}$

$\hookrightarrow Y(s) = 3\frac{e^{-2s}}{s}$

$\hookrightarrow y(t) = \mathcal{L}^{-1}[Y(s)]|_t = \mathcal{L}^{-1}\left[3\frac{e^{-2s}}{s}\right]|_t = 3 \text{step}_2(t) \quad .$

28.6 c. $\mathcal{L}[y'']|_s = \mathcal{L}[\delta_3(t)]|_s$

$\hookrightarrow s^2Y(s) - s \cdot 0 - 0 = e^{-3t}$

$\hookrightarrow Y(s) = \frac{e^{-3s}}{s^2}$

$\hookrightarrow y(t) = \mathcal{L}^{-1}[Y(s)]|_t = \mathcal{L}^{-1}\left[e^{-3s}\frac{1}{s^2}\right]|_t = f(t-3) \text{step}_3(t)$

where

$$f(t) = \mathcal{L}^{-1}\left[\frac{1}{s^2}\right]|_t = t \quad .$$

Thus,

$$f(X) = X \quad , \quad f(t-3) = t-3$$

and

$$y(t) = f(t-3) \text{step}_3(t) = (t-3) \text{step}_3(t) \quad .$$

28.6 e. $\mathcal{L}[y' + 2y]|_s = \mathcal{L}[4\delta_1(t)]|_s$

$\hookrightarrow [sY(s) - 0] + 2Y(s) = 4e^{-s}$

$\hookrightarrow Y(s) = 4e^{-s}\frac{1}{s+2}$

$\hookrightarrow y(t) = \mathcal{L}^{-1}\left[4e^{-s}\frac{1}{s+2}\right]|_t = 4f(t-1) \text{step}_1(t)$

where

$$f(t) = \mathcal{L}^{-1}\left[\frac{1}{s+2}\right]|_t = e^{-2t} \quad .$$

So,

$$f(X) = e^{-2X} \quad , \quad f(t-1) = e^{-2(t-1)}$$

and

$$y(t) = 4f(t-1) \text{step}_1(t) = 4e^{-2(t-1)} \text{step}_1(t) \quad .$$

28.6 g. $\mathcal{L}[y'' + y]|_s = \mathcal{L}[-2\delta_{\pi/2}(t)]|_s$

$\hookrightarrow [s^2Y(s) - s \cdot 0 - 0] + Y(s) = -2e^{-\pi s/2}$

$$\hookrightarrow Y(s) = -2e^{-\pi s/2} \frac{1}{s^2 + 1}$$

$$\hookrightarrow y(t) = \mathcal{L}^{-1} \left[-2e^{-\pi s/2} \frac{1}{s^2 + 1} \right] \Big|_t = -2f\left(t - \frac{\pi}{2}\right) \text{step}_{\pi/2}(t)$$

where

$$f(t) = \mathcal{L}^{-1} \left[\frac{1}{s^2 + 1} \right] \Big|_t = \sin(t) \quad .$$

So,

$$-2f\left(t - \frac{\pi}{2}\right) = -2\sin\left(t - \frac{\pi}{2}\right) = 2\cos(t)$$

and

$$y(t) = -2f\left(t - \frac{\pi}{2}\right) \text{step}_{\pi/2}(t) = \cos(t) \text{step}_{\pi/2}(t) \quad .$$

28.7 a.

$$\mathcal{L}[y' + 3y] \Big|_s = \mathcal{L}[\delta_2(t)] \Big|_s$$

$$\hookrightarrow [sY(s) - \underbrace{y(0)}_2] + 3Y(s) = e^{-2s}$$

$$\hookrightarrow (s + 3)Y(s) - 2 = e^{-2s}$$

$$\hookrightarrow Y(s) = \frac{2}{s + 3} + e^{-2s} \frac{1}{s + 3} \quad .$$

Thus,

$$\begin{aligned} y(t) &= \mathcal{L}^{-1} \left[\frac{2}{s + 3} + e^{-2s} \frac{1}{s + 3} \right] \Big|_t \\ &= 2\mathcal{L}^{-1} \left[\frac{1}{s + 3} \right] \Big|_t + \mathcal{L}^{-1} \left[e^{-2s} \frac{1}{s + 3} \right] \Big|_t = 2f(t) + f(t - 2) \text{step}_2(t) \end{aligned}$$

where

$$f(t) = \mathcal{L}^{-1} \left[\frac{1}{s + 3} \right] \Big|_t = e^{-3t} \quad .$$

So,

$$f(X) = e^{-3X} \quad , \quad f(t - 2) = e^{-3(t-2)}$$

and

$$y(t) = 2f(t) + f(t - 2) \text{step}_2(t) = 2e^{-3t} + e^{-3(t-2)} \text{step}_2(t) \quad .$$

28.7 c.

$$\mathcal{L}[y'' + 3y'] \Big|_s = \mathcal{L}[\delta_1(t)] \Big|_s$$

$$\begin{aligned} \hookrightarrow [s^2 Y(s) - s \underbrace{y(0)}_0 - \underbrace{y'(0)}_1] \\ + 3[sY(s) - \underbrace{y(0)}_0] &= e^{-s} \end{aligned}$$

$$\hookrightarrow (s^2 + 3s)Y(s) - 2 = e^{-s}$$

$$\hookrightarrow Y(s) = \frac{2}{s^2 + 3s} + e^{-s} \frac{1}{s^2 + 3s} \quad . \quad .$$

Thus,

$$\begin{aligned} y(t) &= \mathcal{L}^{-1}\left[\frac{2}{s^2+3s} + e^{-s}\frac{1}{s^2+3s}\right]\Big|_t \\ &= 2\mathcal{L}^{-1}\left[\frac{1}{s^2+3s}\right]\Big|_t + \mathcal{L}^{-1}\left[e^{-1s}\frac{1}{s^2+3s}\right]\Big|_t = 2f(t) + f(t-1)\text{step}_1(t) \end{aligned}$$

where

$$\begin{aligned} f(t) &= \mathcal{L}^{-1}\left[\frac{1}{s^2+3s}\right]\Big|_t = \mathcal{L}^{-1}\left[\frac{1}{s(s+3)}\right]\Big|_t \\ &= \dots \quad (\text{Use partial fractions or convolution.}) \\ &= \frac{1}{3}\left[1 - e^{-3t}\right] \quad . \end{aligned}$$

So,

$$f(X) = \frac{1}{3}\left[1 - e^{-3X}\right] \quad , \quad f(t-1) = \frac{1}{3}\left[1 - e^{-3(t-1)}\right]$$

and

$$y(t) = 2f(t) + f(t-2)\text{step}_2(t) = \frac{2}{3}\left[1 - e^{-3t}\right] + \frac{1}{3}\left[1 - e^{-3(t-1)}\right]\text{step}_1(t) \quad .$$

28.7 e.

$$\mathcal{L}[y'' - 16y]\Big|_s = \mathcal{L}[\delta_{10}(t)]\Big|_s$$

$$\hookrightarrow \left[s^2Y(s) - s \cdot 0 - 0\right] - 16Y(s) = e^{-10s}$$

$$\hookrightarrow Y(s) = e^{-10s} \frac{1}{s^2 + 16}$$

$$\hookrightarrow y(t) = \mathcal{L}^{-1}\left[e^{-10s} \frac{1}{s^2 + 16}\right]\Big|_t = f(t-10)\text{step}_{10}(t)$$

where

$$\begin{aligned} f(t) &= \mathcal{L}^{-1}\left[\frac{1}{s^2+16}\right]\Big|_t = \mathcal{L}^{-1}\left[\frac{1}{(s-4)(s+4)}\right]\Big|_t \\ &= \dots \quad (\text{Use partial fractions or convolution.}) \\ &= \frac{1}{8}\left[e^{4t} - e^{-4t}\right] \quad . \end{aligned}$$

So,

$$f(X) = \frac{1}{8}\left[e^{4X} - e^{-4X}\right] \quad , \quad f(t-10) = \frac{1}{8}\left[e^{4(t-10)} - e^{-4(t-10)}\right]$$

and

$$y(t) = f(t-10)\text{step}_{10}(t) = \frac{1}{8}\left[e^{4(t-10)} - e^{-4(t-10)}\right]\text{step}_{10}(t) \quad .$$

$$28.7 \text{ g.} \quad \mathcal{L}[y'' + 4y' - 12y]|_s = \mathcal{L}[\delta(t)]|_s$$

$$\begin{aligned} \Leftrightarrow \quad & \left[s^2 Y(s) - s \cdot 0 - 0 \right] \\ & + 4[sY(s) - 0] - 12Y(s) = 1 \end{aligned}$$

$$\Leftrightarrow \quad Y(s) = \frac{1}{s^2 + 4s - 12} .$$

So, using either partial fractions or convolution, we find that

$$y(t) = \mathcal{L}^{-1} \left[\frac{1}{s^2 + 4s - 12} \right] \Big|_t = \mathcal{L}^{-1} \left[\frac{1}{(s-2)(s+6)} \right] \Big|_t = \frac{1}{8} [e^{2t} - e^{-6t}] .$$

$$28.7 \text{ i.} \quad \mathcal{L}[y'' + 6y' + 9y]|_s = \mathcal{L}[\delta_4(t)]|_s$$

$$\begin{aligned} \Leftrightarrow \quad & \left[s^2 Y(s) - s \cdot 0 - 0 \right] \\ & + 6[sY(s) - 0] + 9Y(s) = e^{-4s} \end{aligned}$$

$$\Leftrightarrow \quad Y(s) = e^{-4s} \frac{1}{s^2 + 6s + 9}$$

$$\Leftrightarrow \quad y(t) = \mathcal{L} \left[e^{-4s} \frac{1}{s^2 + 6s + 9} \right] \Big|_t = f(t-4) \text{step}_4(t)$$

where

$$\begin{aligned} f(t) &= \mathcal{L}^{-1} \left[\frac{1}{s^2 + 6s + 9} \right] \Big|_t \\ &= \mathcal{L}^{-1} \left[\frac{1}{(s+3)^2} \right] \Big|_t \\ &= \dots \quad (\text{Use the first translation identity or convolution.}) \\ &= te^{-3t} . \end{aligned}$$

So,

$$f(X) = Xe^{-3X} \quad , \quad f(t-4) = (t-4)e^{-3(t-4)}$$

and

$$y(t) = f(t-4) \text{step}_4(t) = (t-4)e^{-3(t-4)} \text{step}_4(t) .$$

$$28.7 \text{ k.} \quad \mathcal{L}[y''' + 9y']|_s = \mathcal{L}[\delta_1(t)]|_s$$

$$\begin{aligned} \Leftrightarrow \quad & \left[s^3 Y(s) - s^2 \cdot 0 - s \cdot 0 - 0 \right] \\ & + 9[sY(s) - 0] = e^{-s} \end{aligned}$$

$$\hookrightarrow Y(s) = e^{-s} \frac{1}{s^3 + 9s}$$

$$\hookrightarrow y(t) = \mathcal{L}^{-1} \left[e^{-s} \frac{1}{s^3 + 9s} \right] \Big|_t = f(t-1) \text{step}_1(t)$$

where

$$\begin{aligned} f(t) &= \mathcal{L}^{-1} \left[\frac{1}{s^3 + 9s} \right] \Big|_t = \mathcal{L}^{-1} \left[\frac{1}{s(s^2 + 9)} \right] \Big|_t \\ &= \dots \quad (\text{Use partial fractions or convolution.}) \\ &= \frac{1}{9} [1 - \cos(3t)] \quad . \end{aligned}$$

So,

$$f(X) = \frac{1}{9} [1 - \cos(3t)] \quad , \quad f(t-1) = \frac{1}{9} [1 - \cos(3[t-1])]$$

and

$$y(t) = f(t-1) \text{step}_1(t) = \frac{1}{9} [1 - \cos(3[t-1])] \text{step}_1(t) \quad .$$